

GATE & PSU's

SAMPLE STUDY MATERIAL

REASONING APTITUDE

GATE & PSU's

Postal Correspondence



CONTENT

DIAGNOSTIC TEST	03-16
1. NUMBER & FRACTIONS	17-34
2. PERCENTAGE	35-49
3. PROFIT AND LOSS	50-66
4. AVERAGE	67-76
5. PERMUTATION AND COMBINATIONS	77-91
6. PROGRESSION	92-109
7. PROBABILITY.....	110-124
8. RATIO AND PROPORTION THEORY	125-129
9. MIXTURES AND ALLIGATION	130-142
10. TIME SPEED & DISTANCES	143-152
11. TIME AND WORK	153-162
12. BOATS AND STREAMS	163-164
13. PIPES AND CISTERN	165-171
14. CALENDAR AND CLOCKS	172-180
15. BLOOD AND RELATIONS	181-187
16. VENN DIAGRAMS	188-193
17. CODING AND DECODING	194-200
18. PIE CHART, BAR CHART BAR DIAGRAM	201-208
19. QUANTATIVE COMPARISON	209-218
20. PUZZLES	219-222
21. REASONING & APTITUDE [MOCK TEST]	223-264
(1) MOCK TEST-1 WITH SOLUTION	223-237
(2) MOCK TEST-2 WITH SOLUTION	238-251
(3) MOCK TEST-3 WITH SOLUTION	252-264

UNIT-1

NUMBER & FRACTIONS

Numeral: In Hindu numeral Arabic system, we can use ten symbols 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 called Digits to represent any number

A group of digits, denoting a number is called a numeral

Place Value

Place value of 2 is $(2 \times 1) = 2$

Place value of 3 is $(3 \times 10) = 30$

Place value of 1 is $(1 \times 100) = 100$ and so on

Place value of 6 is $(6 \times 10^8) = 6000\ 00000$

Face Value: The face value of a digit in a numeral is the value of the digit itself at whatever place. It may be in the above numeral the face value of 2 is 2. The face value of 3 is 3 and so on.

Types of Numbers**1. Natural Numbers**

Counting numbers 1, 2, 3, 4, 5 Are called natural number.

2. Whole Number

All counting number together with zero from the set of whole numbers. Thus

(i) 0 is the only whole number which is not a natural number

(ii) Every natural number is a whole number

3. Integers: All natural numbers, 0 and negatives of counting numbers $(-3, -2, -1, 0, 1, 2, 3 \dots\dots)$ together from the set of integers

1. Positive integers 2. Negative integers

3. Non positive and non positive integers

4. Even Numbers

A number divisible by 2 is called an even numbers

5. Odd Number

A number not divisible by 2 is called an odd number 1, 3, 5, 7, 9.

6. Prime Number

A number greater than 1 is called prime number if it is divisible by either 1 or it self.

Prime number up to 100 are 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 83, 89, 97

(ii) Prime numbers greater than : Let p be given number greater than 100. To find out whether it is prime or not we use the following method.

(iii) Find a whole number nearly greater than the square root of p .

(iv) Prime numbers less than 14 are 2, 3, 5, 7, 11, 13

(v) 191 is not divisible by any of them. So, 191 is a prime number.

Composite Number

Numbers greater than 1 which are not prime are known as composite no., e.g. 4, 6, 8, 9, 10, 12.

(i) 1 is neither prime nor composite

(ii) 2 is the only even number which is prime

(iii) There are 25 prime numbers between 1 and 100

Test of Divisibility

1. Divisibility by 2

A number is divisible by 2. If its unit's digit is any of 0, 2, 4, 6, 8.

2. Divisibility by 3

A number is divisible by 3. If the sum of its digits is divisible by 3.

3. Divisibility by 4

A number is divisible by 4. If the number formed by the last two digits is divisible by 4.

4. Divisibility by 5

A number is divisible by 5. If its unit's digit is either 0 or 5. Thus, 20820 and 50345 are divisible by 5. While 30934 and 40946 are not.

Divisibility by 6

A number is divisible by 6. If its is divisible by both 2 and 3.

Divisibility by 8

A number is divisible by 8. If the number formed by the last three digits of the given number is divisible by 8.

Divisibility by 9

A number is divisible by 9. If the sum of its digits is divisible by 9.

(A) Multiplication by short cut methods

(i) $a * (b + c) = a \times b + a \times c$

(ii) $a(b - c) = ab - ac$

(B) Multiplication of a number by 5^n

Put n zeroes to the right of the multiplicand and divide. The no. so formed divided by 2^n

Division Algorithm or Euclidean Algorithm

If we divide A given number by another number. Then dividend = (Divisors \times Quotient) + Remainder

Progression

Arithmetic Progression

N th term of this A.P.

$$T_n = a + (n - 1) d$$

$$S_n = \frac{n}{2}[2a + (n-1)d] = \frac{n}{2}[\text{first term} + \text{last term}]$$

$$(A) (1 + 2 + 3 + \dots + n) = \frac{n(n+1)}{2} \quad (B) (1^2 + 2^2 + 3^2 + \dots + n^2) = \frac{n(n+1)(2n+1)}{6}$$

$$(C) (1^3 + 2^3 + \dots + n^3) = n^2 \frac{(n+1)^2}{4}$$

Geometrical Progression

In this G.P. = $T_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{(1-r)} \text{ when } r < 1$$

$$S_n = \frac{a(r^n-1)}{r-1} \text{ when } r > 1$$

Decimal Fractions

Fractions in which denominator are powers of 10 are known as decimal fractions

$$\frac{1}{100} = 1 \text{ hundredth} = .01$$

Conversion of a Decimal into Vulgar fractions

Put 1 in the denominator under the decimal point and annex with it as many zeroes as is the number of digits. After the decimal point. Now remove the decimal point and reduce the fraction to its lowest terms

Thus $0.25 = \frac{25}{100} = \frac{1}{4}$

Recurring Decimal

If in a decimal fraction, A figure or a set of figures is repeated continuously. Then such A number is called A recurring decimal. In a recurring decimal, If a single figure is repeated. Then it is expressed by putting A dot on it. If a set of figures is repeated it is expressed by putting A bar on the set

$$\frac{1}{3} = 0.3333 = 0.\dot{3}$$

$$\frac{22}{7} = 3.142857142857 \dots\dots\dots$$

Types of Recurring Decimal

- (1) Pure Recurring Decimal
- (2) Mixed Recurring Decimal

(1) Pure Recurring Decimal

A decimal fraction in which all the figures after the decimal point are repeated, is called. A pure receding decimal.

Example: $0.\bar{5} = \frac{5}{9}$; $0.\bar{53} = \frac{53}{99}$, $0.\overline{067} = \frac{67}{999}$ etc.

(2) Mixed Recurring Decimal

A decimal fraction in which some figures do not repeat and some of them are repeated is called a mixed recurring decimal.

Example: $0.17\overline{33} = \frac{1733-17}{9900} = 1716/9900$

Some Examples:

1. If $a + \frac{1}{1 + \frac{1}{3 + \frac{1}{4}}} = 2$ then $a = ?$

Solution:

$$a + \frac{1}{1 + \frac{1}{3 + \frac{1}{4}}} = 2, \quad a + \frac{1}{1 + \frac{1}{(13/4)}} = 2 \Rightarrow a + \frac{1}{1 + \frac{4}{13}} = 2 \Rightarrow a + \frac{1}{17/13} = 2$$

$$\Rightarrow a + \frac{13}{17} = 2 \Rightarrow a = 2 - \frac{13}{17} \Rightarrow a = \frac{34-13}{17} = \frac{21}{17} = 1\frac{4}{17}$$

2. On children's day sweets were to be equally distributed amongst 540 children. But on that particular day, 120 children were absent. Thus, each child got 4 sweets extra. How many sweets was each originally supposed to get?

Solution: Suppose, each child was supposed to get x sweets. Then we have,

$$540 \times x = (540 - 120)(x + 4)$$

$$\Rightarrow 120x = 420 \times 4 \Rightarrow x = \frac{420 \times 4}{120} = 14$$

Therefore, $x = 14$ sweets.

3. The difference between the squares of two consecutive odd integers is always divisible by which digits or numbers?

Solution: Let the two consecutive odd numbers be $(2n + 1)$ and $(2n + 3)$ respectively.

Then,

$$(2n + 3)^2 - (2n + 1)^2 = (2n + 3 + 2n + 1)(2n + 3 - 2n - 1)$$

$$= (4n + 4) \times 2 = 8(n + 1), \text{ which is divisible by } 8.$$

4. In an entrance examination of GATE, there are two sections of Biological science students and the two sections are Life sciences and Biotechnology. If 10 students of Biotechnology shift over to Life science, the strength of appearing students in Life science becomes three times the strength of Biotechnology. But, if 10 students shift over from Life science to Biotechnology, both Life science and Biotechnology students becomes equal. How many students there in Life Science and Biotechnology?

- (a) 50 and 30 (b) 45 and 15 (c) 90 and 40 (d) 30 and 50

ANS: a

5. Simplify the given fraction: $12 \div \frac{1}{7 - \frac{1}{1 - \frac{1}{1 + \frac{1}{2}}}}$ of $19\frac{1}{5}$

- (a) 2/5 (b) 3/4 (c) 5/2 (d) 4/3

ANS: c

6. "October2, 2001" in MMDDYYYY format is a palindrome (a string that reads the same forwards as it does backwards example, 10/02/2001.....10022001). When was the latest century before October 2, 2001 that is also a palindrome?

- (a) 13th century (b) 14th century (c) 17th century (d) 20th century

ANS: b

7. Find the largest natural number which exactly divides the product of any 4 consecutive natural numbers.

Solution: Required number = $1 \times 2 \times 3 \times 4 = 24$

Therefore, required number = 24 [it is applicable for all 4 consecutive natural numbers]

8. Murari, Arun and Nitin start at the same time in the same direction to run around a circular stadium. Murari completes a round in 126 seconds, Arun in 154 seconds and Nitin in 99 seconds, all starting at the same point. After what time they meet again at the starting point?

Solution: Required time = L.C. M. of 126 sec, 154 sec and 99 sec.

$$\begin{array}{r} 2 \mid 126, 154, 99 \\ 11 \mid 63, 77, 99 \\ 9 \mid 63, 7, 9 \\ 7 \mid 7, 7, 1 \\ \hline 1, 1, 1 \end{array}$$

Therefore, required time = $2 \times 110 \times 9 \times 7 = 1386$ sec = 23 min. 10 sec.

9. Amit, Bijender and Chandu go walking round a circle 1 km in circumference at the rates of 10 m/min, 20 m/min and 40m/min respectively, if they all start together and walk in the same direction, when will they be together at the same place?

- (a) After 50 minute (b) after 100 minute
(c) after 240 minute (d) after 800 minute

ANS: b

10. In an examination, a student average marks were 63 per paper. If he had obtained 20 more marks for his Biochemistry paper and 2 more marks for his Microbiology paper, his average per paper would have been 65. How many papers were there in the examinations?

Solution: Assume there by x all papers.

Total marks of all papers = 63x

From question

$$65x - 63x = 20 + 2$$

$$\Rightarrow 2x = 22 \quad \therefore x = 11$$

11. The work done by a child is one third that by a man and half that by a woman. If one man, one woman and one child together can complete a work in 2 days, in how many days can 4 children together complete the same work?

Solution: We have, $1M = 3C$ and $1W = 2C$ (where, M = male work, C = child work, W = woman work)

Therefore, $1M + 1W + 1C = 6C$

Hence, the required number of days = $\frac{6 \times 2}{4} = 3 \text{ days}$

12. In a central government office, there are 5 working days and for each day, the working hours are 8. An employee gets Rs. 2.40 per hour for regular work and Rs. 3.20 per hour for overtime. If he earns Rs. 432 in 4 weeks, how many hours he work for?

(a) 195 (b) 160 (c) 175 (d) 180

ANS: c

13. Nitin and Pradip solved a quadratic equation. In solving it, Nitin made a mistake in the constant term and got the roots as 6 and 2, while Pradip made a mistake in the coefficient of x only and obtained the roots as -7 and -1. Find the correct roots of equation

Solution: For Nitin, we have $r + s = 8$ and $rs = 12$

The equation is $x^2 - (\text{sum of roots})x + \text{product of roots} = 0$

$$\therefore x^2 - 8x + 12 = 0$$

For Pradip, we have $r + s = -8$ and $rs = 7$

Therefore, the equation is $x^2 + 8x + 7 = 0$

When there is no mistake in a and b, the sum of roots must be correct.

Therefore, sum of roots = $6 + 2 = 8$ and product of roots = $(-7) \times (-1) = 7$

So, the correct equation is $x^2 - 8x + 7 = 0$


$$\Rightarrow x^2 - 7x - x + 7 = 0$$

$$\Rightarrow (x - 7)(x - 1) = 0$$

$$\Rightarrow x = 7 \text{ or } x = 1$$

Therefore, the roots are 7, 1.

14. Pradeep wishes to make a gravel path around his rectangular pond. The path must be the same width all the way round, as shown in the diagram. The pond measures 4m by 9m and he has enough gravel to cover an area of 48m^2 . How wide around the path be?



- (a) 2 meter (b) 3 meter (c) 1.5 meter (d) 8 meter

ANS: c

15. A man arranges to pay off a debt of Rs. 3600 by 40 annual installments which are in A.P. when 30 of the installments are paid, the dues leaving one third of the debt unpaid. Find the value of 8th installment.

Solution: Let the first installment be a and common difference of A.P be d .

$$\text{Given, } 3600 = \text{sum of 40 terms} = \frac{40}{2} \{2a + (40 - 1)d\}$$

$$\Rightarrow 3600 = 20\{2a + 39d\}$$

$$\Rightarrow 2a + 39d = 180 \quad \dots(i)$$

After 30 installments, one third of the debt is unpaid

$$\text{Hence, } \frac{3600}{3} = 1200 \text{ is unpaired and } 2400 \text{ is paid.}$$

$$\text{Now, } 2400 = \frac{30}{2} \{2a + (30 - 1)d\}$$

$$\Rightarrow 2400 = 15\{2a + 29d\}$$

$$\Rightarrow 2a + 29d = 160 \quad \dots(ii)$$

Subtracting (ii) from (i), we get

$$\Rightarrow 10d = 20 \quad \therefore d = 2$$

$$\text{From (i), } 2a = 180 - 39d = 180 - 39 \times 2 = 180 - 78 = 102$$

$$\text{Therefore, } a = 51$$

$$\text{Now value of the 8th installment} = a + (8 - 1)d = 51 + 7 \times 2 = 65$$

16. The population of bacteria culture doubles every 2 minutes. How many minutes will it takes for the population grow from 1000 to 512000 bacteria?

Solution: Let the growth be 2000, 4000,512000

This is the G.P in which $a = 2000$, $r = 2$ and $t_n = 512000$

$$\text{Since, } t_n = ar^{n-1}$$

$$\Rightarrow 512000 = 2000 \times 2^{n-1}$$

$$\Rightarrow 512 = 2^{n-1} \times 2^1$$

$$\Rightarrow 2^n = 2^9$$

$$\Rightarrow n = 9$$

Therefore, time taken = $2 \times 9 = 18$ minutes.

17. If $\log_{\sqrt{8}} x = 3\frac{1}{3}$ then $x = ?$

Solution: $\log_{\sqrt{8}} x = 3\frac{1}{3} = \frac{10}{3}$

$$\Rightarrow x = (\sqrt{8})^{\frac{10}{3}} = (2^{3/2})^{\frac{10}{3}} = 2^{\left(\frac{3 \times 10}{3}\right)} = 2^5 = 32 \quad \therefore x = 32$$

18. If $\log_{10} 2 = 0.3010$ and $\log_{10} 3 = 0.4714$, find the value of $\log_{10} 25$ and $\log_{10} 4.5$.

Solution: $\log_{10} 25 = \log_{10} \left(\frac{100}{4}\right) = \log_{10} 100 - \log_{10} 4 = \log_{10} 10^2 - \log_{10} 2^2$


$$= 2 - 2\log_{10} 2 = 2 - 2 \times 0.3010 = 1.398$$

$$\log_{10} 4.5 = \log_{10} \left(\frac{9}{2}\right) = \log_{10} 9 - \log_{10} 2 = \log_{10} 3^2 - \log_{10} 2$$

$$= 2\log_{10} 3 - \log_{10} 2 = 2 \times 0.4771 - 0.3010 = 0.6532$$

19. What is the minimum value of the perimeter of a triangle, if two of its sides are 5 cm and 7 cm respectively? (the sides have integer volumes)

Solution: In a triangle, $c > b - a$




Therefore, $c > 7 - 5$ or $c > 2$

The minimum length of the third side is 3 cm

Hence, the minimum value of the perimeter = $5 + 7 + 3 = 15$ cm

20. What is the area of rectangle EFGH given figure, if the area of rectangle ABCD is 100?



Solution. Area of rectangle ABCD $= (x+1) \times (x+4) = x^2 + 5x + 4$

Area of rectangular EFGH $= (x+2) \times (x+3) = x^2 + 5x + 6$

If we compare the area of rectangle ABCD and EFGH, the area of rectangle EFGH is 2 more than the area of rectangle ABCD.

Therefore, area of rectangle EFGH $= 100 + 2 = 102$

LEVEL-1

1. Which of the following are prime number (i) 241 (ii) 337 (iii) 391
(a) 241, 337 (b) 337, 391 (c) All the above (d) None of these
2. Find the unit's digit in $(264)^{102} + (264)^{103}$
(a) 0 (b) 2 (c) 4 (d) 20
3. Find the total number of prime factors in the expression $(4)^{14} \times (7)^5 \times (11)^2$
(a) 1 (b) 2 (c) 3 (d) 4
4. Simplify (i) $896 \times 896 - 204 \times 204$ (ii) $387 \times 387 + 114 \times 114 + 2 \times 387 \times 114$
(a) 761000, 251000 (b) 761200, 251001 (c) 761200, 251000 (d) 76100, 25100
5. Which of the following numbers is divisible by 3 (i) 541326 (ii) 5967013
(a) (i) and (ii) (b) only (ii) (c) only (d) None of these
6. Which one of the following is not a prime number
(a) 31 (b) 61 (c) 71 (d) 91
7. $(112 * 5^4) = ?$
(a) 67000 (b) 50055 (c) 70000 (d) 75000
8. What least value must be assigned * so that the number $54623*7$ is exactly divisible by 9
(a) 1 (b) 0 (c) 2 (d) 3
9. Which of the following numbers is divisible by 4
(i) 6792059 (ii) 618703572
(a) (i) and (ii) (b) only (ii) (c) only (i) (d) None of these
10. Which digits should come in place of * and \$. If the number $62684 * \$$ is divisible by both 8 and 5.
(a) 4, 0 (b) 0, 4 (c) 4, 4 (d) none of these
11. Is 4832718 is divisible by 11
(a) Yes (b) No (c) Data inadequate (d) None of these
12. A number when divided by 342 gives a remainder 47. When the same number is divided by 19, what would be the remainder
(a) 5 (b) 9 (c) 4 (d) 0
13. Find the remainder when 2^{31} is divided by 5
(a) 4 (b) 5 (c) 3 (d) 7

14. Find the sum of all odd numbers up to 100
 (a) 2000 (b) 2500 (c) 2800 (d) 3000
15. Find the sum of all 2 digit numbers divisible by 3
 (a) 1700 (b) 1665 (c) 1600 (d) 1605
16. It is being given that $(2^{32} + 1)$ is completely divisible by a whole number which of the following numbers is completely divisible by this number.
 (a) $(2^{16} + 1)$ (b) $(2^{16} - 1)$ (c) $7 * 2^{23}$ (d) $2^{96} + 1$
17. How many prime numbers are less than 50?
 (a) 16 (b) 15 (c) 14 (d) 18

BRIEF SOLUTIONS

1. 391 is not prime number , 337 is a prime number, 241 is a prime number

2. Required unit's digit = unit digit in $(4)^{102} + (4)^{103}$

Now 4^2 gives unit digit 6

$\therefore (4)^{102}$ gives unit digit 6

$\therefore (4)^{103}$ gives unit digit of the product $(6 \times 4) = 4$

Hence unit's digit in $(264)^m + (264)^{103} = \text{unit digit in } (6 + 4) = 0$

3. Total no. of prime factors = number of bases = 3

4. $(896)^2 - (204)^2$
 $= (896 + 204)(896 - 204) = 1100 \times 692 = 761200$

(ii) Given expression

$$= (387)^2 + (114)^2 + (2 \times 387 \times 114) = a^2 + b^2 + 2ab$$

where $a = 387$ $b = 114$ $= (501)^2 = 251001$

5. Sum of digits in 541326 = 21
 Which is divisible by 3 hence 541326 is divisible by 3.
 Sum of digits in 5967013 = 31
 Which is not divisible by 3, hence 5967013 is not divisible by 3.
6. 91 is divisible by 7, so it is a prime number.
7. 70000

8. Ans. (b)

9. (b)

10. (4, 0)

11. (Sum of digits at odd places) – (Sum of digits at even places)
 $= (8 + 7 + 3 + 4) - (1 + 2 + 8) = 11$

Which is divisible by 11. Hence 4832718 is divisible by 11.

12. On dividing the given number by 342, Let k be the quotient and 47 as remainder.

Then number $= 342k + 47 = 19(18k + 2) + 9$

\therefore The given no when divided by 19 gives $(18k + 2)$ as quotient and 9 as remainder

13. $2^{10} = 1024$ unit digit of $2^{10} \times 2^{10} \times 2^{10}$ is 4

as $4 \times 4 \times 4$ gives unit digit 4

\therefore Unit digit of 231 is 8. Now 8 when dividend by 5, gives 3 as remainder.

Hence 231 when dividend by 3, gives as remainder.

14. The given numbers are 1, 3, 5, 7, 99

This is an A.P. with $a = 1$ and $d = 2$

Let it contains n terms

$$= \text{Then } 1 + (n - 1) \times 2 = 99 \quad \text{or} \quad n = 50$$

$$\therefore \text{ Required sum} = n(\text{first term} + \text{last term}) = \frac{50}{2}(1 + 99) = 2500$$

15. Required sum $= \frac{30}{2}(12 + 99) = 1665$

16. Let

$$2^{32} = x$$

Then $(2^{32} + 1) = (x + 1)$

Let $(x + 1)$ be completely divisible by the natural number N. Then $(2^{96} + 1) = [(2^{32})^3 + 1]$

$$= (x^3 + 1) = (x + 1)(x^2 - x + 1)$$

Which is completely divisible by N.

Since $(x + 1)$ is divisible by N.

17. Prime numbers less than 50 are
2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 47 their no. is 14.

LEVEL-2

1.
$$\frac{(489 + 375)^2 - (489 - 375)^2}{(489 \times 375)}$$

(a) 144 (b) 864 (c) 2 (d) 4
2. Given that $268 \times 74 = 19832$. Find the values find the values of 2.68×0.74
(a) 198.32 (b) 19.832 (c) 1.9832 (d) 1983.2
3. Find the quotient $\frac{0.0204}{17}$
(a) 0.0012 (b) 0.012 (c) 0.00012 (d) None of these
4. What value will come in place of question mark in the following question $0.0006 + ? = 0.6$
(a) 0.01 (b) 0.1 (c) 0.001 (d) 0.0001
5. If $\frac{1}{3.718} = 0.2689$. Then find the value of $\frac{1}{0.0003718}$
(a) 2698 (b) 2698 (c) 2968 (d) 9286
6. Express as Vulgar fractions $0.\overline{37}$
(a) $\frac{99}{37}$ (b) $\frac{0.37}{99}$ (c) $\frac{37}{99}$ (d) $\frac{99}{0.37}$
7. Express as Vulgar fractions $0.\overline{053}$
(a) $\frac{99}{0.37}$ (b) $\frac{53}{999}$ (c) $\frac{999}{0.53}$ (d) $\frac{0.53}{999}$
8. Simplify $\frac{0.05 \times 0.05 \times 0.05 + 0.04 \times 0.04 \times 0.04}{0.05 \times 0.05 - 0.05 \times 0.04 + 0.04 \times 0.04}$
(a) 0.9 (b) 0.09 (c) 9 (d) 0.009
9. Find the product of $0.4 \times 0.04 \times 0.004 \times 40$
(a) 0.002650 (b) 0.002560 (c) 0.02560 (d) 0.05260
10. How many digits will be there to the right of the decimal point in the product of 95.75 and 0.2554.
(a) 5 (b) 6 (c) 7 (d) None of these
11. Which of the following is equal is 3.14×10^6
(a) 314 (b) 314000 (c) 3140 (d) None of these
12. $|P - 10| = 12$ & $|4J - 10| = 6$
What is the max value of P/J

- (a) 11 (b) 22 (c) -2 (d) None of these
13. If $\sqrt{a^b} = 5b + a^2$ then (a, b) could be
 (a) (3, 4) (b) (2, 12) (c) (4, 18) (d) (6, 4)
14. The value of $\sqrt{6 + \sqrt{6 + \sqrt{6 + \sqrt{6 + \dots}}}}$ is
 (a) 2 (b) 5 (c) 4 (d) 3
15. Find the value of
 $(28 + 10\sqrt{3})^{1/2} - (7 - 4\sqrt{3})^{1/2}$
 (a) 5 (b) 3 (c) 8 (d) 6

BRIEF SOLUTIONS

1. Given expression = $\frac{(a+b)^2 - (a-b)^2}{ab} = 4$
2. **Ans. c**
3. $\frac{204}{17} = 12$ dividend contains 4 places of decimal $\frac{0.2040}{17} = 0.0012$
4. Let $\frac{0.006}{x} = 0.6$
 Then $x = 0.01$
5. $\frac{1}{0.0003718} = 2689$
6. $0.\overline{37} = \frac{37}{99}$
7. $0.\overline{053} = \frac{53}{999}$
8. 0.09
9. Sum of decimal places = $(1 + 2 + 3) = 6$
 So $0.4 \times 0.04 \times 0.004 \times 40 = 0.002560$
10. Sum of decimal digits = 7
11. 3140000
12. (b)
 max \Rightarrow P max, T min
 $|P - 10| = 12 \Rightarrow P = 22 \text{ or } -2$
 $|4T - 10| = 6 \Rightarrow T = 4 \text{ or } 1$
 So max value of 22.
13. (b) **Put** $a = 2, b = 12$
 $2^6 = 64$

14. (d) Let $x = \sqrt{6 + \sqrt{6 + \sqrt{6 + \dots}}}$

$$\Rightarrow x = \sqrt{6 + x}$$

$$\Rightarrow x^2 = 6 + x$$

$$\Rightarrow x^2 - x - 6 = 0$$

$$\Rightarrow (x - 3)(x + 2) = 0$$

$$\Rightarrow x = 3$$

15. (b)

$$(28 + 10\sqrt{3})^{1/2} - (7 - 4\sqrt{3})^{-1/2}$$

$$= (5 + \sqrt{3}) - (2 - \sqrt{3})^{-1}$$

$$\Rightarrow (5 + \sqrt{3}) - \frac{1}{2 - \sqrt{3}} \times \frac{2 + \sqrt{3}}{2 + \sqrt{3}}$$

$$\Rightarrow (5 + \sqrt{3}) - (2 + \sqrt{3}) = 3$$

SIMPLIFICATION OF ROOTS

Prerequisites

(1) BODMAS Rule

B → Bracket

O → Of

D → Division

M → Multiplication

A → Addition

S → Subtract

Thus in simplifying an expression first of all the brackets must be removed, strictly in the order

(), { }, | |

After removing the brackets

(i) of (ii) division

(iii) multiplication

(iv) subtraction

Modules of a Real Number

$$|a| = \begin{cases} a & \text{If } a > 0 \\ -a & \text{If } a < 0 \end{cases}$$

Virnaculum: (or Bar)

When an expression contains virnaculum before applying the 'BODMAS' Rule

Roots

Roots or (Radicals) are the 'opposite' operation of applying exponents you can 'undo' A power with a radical and A radical can 'undo' A power.

For instance, If you square 3, you get 9, and if you 'take the square root?' you get 3.

$2^2 = 4$

$\sqrt{4} = 2$

$3^2 = 9$

$\sqrt{9} = 3$

LEVEL-3

1. A man has Rs. 480 in the denominations of one rupee notes, five rupee notes and ten rupee notes the number of notes of each denomination is equal. What is the total no. of notes

.....Sample file



**To Buy Postal Correspondence Package call at
0-9990657855**